

On Detecting Discrete Cheshire Charge[†]

Martin Bucher, Kai-Ming Lee, and John Preskill

California Institute of Technology, Pasadena, CA 91125

Abstract

We analyze the charges carried by loops of string in models with non-abelian local discrete symmetry. The charge on a loop has no localized source, but can be detected by means of the Aharonov–Bohm interaction of the loop with another string. We describe the process of charge detection, and the transfer of charge between point particles and string loops, in terms of gauge-invariant correlation functions.

[†] This work supported in part by the US Department of Energy under Contract No. DE-AC03-81-ER40050

1. Introduction

In a spontaneously broken gauge theory, if the unbroken gauge group H is a *discrete* subgroup of the underlying continuous gauge group G , then the theory will contain topologically stable strings (in 3+1 dimensions) or vortices (in 2+1 dimensions). If H is non-abelian, the strings have remarkable properties. In particular, a closed loop of string can carry a nontrivial H charge. Oddly, this charge is a global property of the string that can not be attributed to any locally defined charge density. Yet the charge is physically detectable, for the charged string loop has an infinite range Aharonov–Bohm interaction with other strings. Furthermore, if a pointlike particle carrying H charge winds through a string loop, the particle and the loop can exchange charge.

Charge with no localized source has been called “Cheshire charge.”^[1] It was first discussed for the case of the “Alice” string.^[2] A loop of Alice string can carry electric charge, and have a long-range electric field, even though the electric charge density vanishes everywhere.^[3,1,4] Processes in which electric (or magnetic) charge is exchanged between string loops and point particles were discussed in Ref. 1, 4-6.

In this paper, we analyze the purely quantum-mechanical version of Cheshire charge that arises in a theory with a non-abelian discrete local H symmetry.^[7,4] The semiclassical theory of discrete Cheshire charge was formulated in Ref. 4, and elaborated in Ref. 8, 9. Here we extend the theory further, in several respects. We describe how a charge operator can be constructed, such that the expectation value of the operator in a state specifies the transformation properties of the state under global H transformations. We then study processes in which charge is exchanged between string loops and point particles, and derive general formulas for how the expectation value of the charge of the loop is altered by the exchange. Finally, we explain how the charge exchange processes can be probed using gauge-invariant correlation functions.

The charge operator and correlation functions are also treated in Ref. 10, where lattice realizations of operators and correlators are extensively discussed.

The rest of this paper is organized as follows: In Section 2, we briefly review the basic properties of non-abelian strings and the concept of Cheshire charge. We construct the non-abelian charge operator in Section 3, and analyze the charge exchange process in Section 4. Section 5 contains a final comment.

2. Non-Abelian Strings

Let us briefly recall some of the properties of non-abelian strings in three spatial dimensions (and vortices in two spatial dimensions).

If a simply connected gauge group G is broken to a discrete subgroup H , then strings are classified by elements of H . To assign a group element to a loop of string, we fix an (arbitrary) basepoint x_0 , and specify a path C , beginning and ending at x_0 , that winds once through the string loop. (See Fig. 1.) The assigned group element is then

$$a(C, x_0) = P \exp \left(i \int_{C, x_0} A \cdot dx \right) . \quad (2.1)$$

We refer to $a(C, x_0)$ as the “flux” of the string; it encodes the effect of parallel transport around the path C . The flux takes values in $H(x_0)$, the subgroup of G that stabilizes the Higgs condensate at the point x_0 (since parallel transport around C must return the condensate to its original value). Since H is discrete, the flux $a(C, x_0)$ is unchanged by deformations of C that leave x_0 fixed, as long as C never crosses the core of the string.

For a configuration of many string loops, we specify a standard path for each loop, where all paths have the same basepoint. Evidently, the flux associated with the product path $C_2 \circ C_1$ obtained by traversing first C_1 and then C_2 is just the product $a(C_2, x_0) \cdot a(C_1, x_0)$ of the two fluxes associated with C_1 and C_2 . Thus, $a(C, x_0)$ defines a homomorphism that maps $\pi_1(\mathcal{M}, x_0)$ to H , where $\mathcal{M}$ is the manifold that is obtained when the cores of all strings are removed from $\mathbb{R}^3$.

The flux assigned to a path is not gauge invariant. The gauge transformations at the basepoint x_0 that preserve the condensate at the basepoint, and so preserve the embedding of H in G , take values in $H(x_0)$. Under such a gauge transformation $h \in H(x_0)$, the flux transforms as

$$a(C, x_0) \rightarrow h a(C, x_0) h^{-1} . \quad (2.2)$$

In a many-string configuration, the flux of each string becomes conjugated by h .

In the presence of strings, the embedding of the unbroken group H in G necessarily depends on the spatial position x . If the strings are non-abelian, this position dependence is described by a nontrivial fiber bundle. The base space of the bundle is the spatial manifold $\mathcal{M}$, the fiber is H , and the structure group is also H , which acts on the fiber by conjugation. The bundle is twisted: Upon transport around the path C , the group element $h \in H(x_0)$ becomes conjugated by $a(C, x_0)$. This twist prevents the bundle from being smoothly deformed to the trivial bundle $\mathcal{M} \times H$. One thus says that the unbroken H symmetry is not “globally realizable;”^[11,1,4] there is no smooth function of position that describes how the unbroken group is embedded in G . Only the subgroup of H that commutes with the flux of all strings is globally realizable on $\mathcal{M}$.

To define the H -charge of a state, we will want to consider how the state transforms under global H transformations. Fortunately, these global gauge transformations can be implemented, even though a topological obstruction prevents H from being globally realized. The point is that it is sufficient to be able to define an H transformation on and outside a large surface Σ (homeomorphic to S^2) that encloses all of the string loops. The transformation cannot be smoothly extended inside the sphere if it is required to take values in $H(x)$. However, one may relax this requirement and allow the gauge transformation to take values in G inside of Σ ; then a smooth extension is possible. It makes no difference what extension is chosen, for gauge transformations of compact support act trivially on physical states. (In two spatial dimensions, the only global H transformations that can be implemented are

those that commute with the *total* flux; *i.e.*, the flux associated with a path that encloses all of the vortices.)

If the basepoint x_0 lies outside the surface Σ , then, under the global gauge transformation $h \in H$, the flux of a string transforms as in eq. (2.2). Thus, the H representations mix up the string loop state labeled by $a \in H$ with string loop states labeled by other group elements in the same conjugacy class as a . Let $[a]$ denote the conjugacy class that contains a . The action of H on the members of the class $[a]$ defines a (reducible) representation that we denote as $D^{([a])}$. In $D^{([a])}$, each element of H is represented by a permutation of the class, according to

$$D^{([a])}(h) : |a'\rangle \rightarrow |ha'h^{-1}\rangle, \quad a' \in [a]. \quad (2.3)$$

This representation can be decomposed into irreducible representations of H . For each class $[a]$ there is a unique state that can be constructed that transforms trivially under H ; it is the superposition of flux eigenstates

$$|0; [a]\rangle = \frac{1}{\sqrt{n_{[a]}}} \sum_{a' \in [a]} |a'\rangle, \quad (2.4)$$

where $n_{[a]}$ denotes the order of the class. The other states contained in the decomposition of $D^{([a])}$ carry H -charge. This is “discrete Cheshire charge,” for the charge of the loop has no localized source. (Note that the charged string states transform trivially under the *center* of H , since $D^{([a])}$ represents the center trivially.)

The splitting between the charge-0 string state eq. (2.4) and the lowest charge excitation of the string is of order $\exp(-\kappa A)$, where κ is a string tension, and A is the area of the string loop.^[9,10] It is a remarkable property of Cheshire charge that, in the presence of a large string loop, the gap between the ground state and the first charged excitation is much less than the corresponding gap when the string is absent. Indeed, the gap approaches zero very rapidly as the size of the loop increases.

3. Charge Operator

The discrete charge of an object, including a charged string loop, can be detected at long range by means of the Aharonov–Bohm effect.^[12] Let $|u\rangle$ denote the wave-function in internal–symmetry space of an object located at x_0 that transforms as the irreducible representation $D^{(\nu)}$ of H . Then when the particle is transported around the closed path C that begins and ends at x_0 , the wave-function is modified according to

$$|u\rangle \rightarrow D^{(\nu)}[a(C, x_0)] |u\rangle ; \quad (3.1)$$

if the string is in the flux eigenstate $|a\rangle$, then the Aharonov–Bohm phase that can be measured in an interference experiment is

$$\langle u| D^{(\nu)}(a) |u\rangle . \quad (3.2)$$

But if the string is in the charge–zero eigenstate $|0; [a]\rangle$ given by eq. (2.4), then the expectation value of the “phase” $D^{(\nu)}(a)$ becomes

$$\frac{1}{n_{[a]}} \sum_{a' \in [a]} D^{(\nu)}(a') = \frac{1}{n_H} \sum_{h \in H} D^{(\nu)}(hah^{-1}) = \frac{1}{n_\nu} \chi^{(\nu)}(a) \mathbf{1} , \quad (3.3)$$

where n_H is the order of the group, n_ν is the dimension of $D^{(\nu)}$, and $\chi^{(\nu)}$ is the character of the representation. The second equality follows from Schur’s lemma.

In principle, the charge inside a large region can be measured by means of a process in which the world sheet of a string sweeps over the boundary of the region. If the string is in the charge-zero eigenstate $|0; [a]\rangle$, and the object enclosed by the world sheet transforms as the irreducible representation (ν) of H , then the amplitude for this process will be weighted by the Aharonov–Bohm factor $(1/n_\nu)\chi^{(\nu)}(a)$. The charge (ν) of an unidentified object can be determined by measuring this factor for each class $[a]$.

A gauge-invariant operator $F_{[a]}(\Sigma)$ can be constructed that inserts, as a classical source, a string worldsheet in the state $|0; [a]\rangle$ on the closed surface Σ . The realization of this operator in a Euclidean lattice gauge theory was described in Ref. 4 in the case where H is abelian (see also Ref. 13), and in Ref. 14, 10 for H non-abelian. (It is closely related to the 't Hooft loop operator.^[15]) If the surface Σ is chosen to lie in a time slice, then the operator $F_{[a]}(\Sigma)$ measures the charge enclosed by Σ . To define the charge of an isolated object, we consider a surface Σ that encloses the object, and whose closest approach to the object is large compared to the correlation length of the theory. Let $|\psi\rangle$ denote the quantum state of the object. Then we have

$$\frac{\langle\psi| F_{[a]}(\Sigma) |\psi\rangle}{\langle F_{[a]}(\Sigma)\rangle_0} = \sum_{\nu} p^{(\nu)}(\psi; \Sigma) \frac{1}{n_{\nu}} \chi^{(\nu)}(a) , \quad (3.4)$$

where $p^{(\nu)}(\psi; \Sigma)$ is the probability that the object carries charge (ν) . By measuring $F_{[a]}(\Sigma)$ for each class, we can determine all of the $p^{(\nu)}$'s. (It is necessary to divide by the vacuum expectation value $\langle F_{[a]}(\Sigma)\rangle_0$ to remove the effects of quantum-mechanical vacuum charge fluctuations near the surface Σ .^[4])

The Aharonov-Bohm interaction makes it possible to detect H -charge at arbitrarily long range. Thus, a theory with discrete local H symmetry obeys a charge superselection rule—no gauge-invariant local operator can create or destroy H -charge. We have

$$\langle(\mu)| \mathcal{O} |(\nu)\rangle = 0 , \quad (\mu) \neq (\nu) , \quad (3.5)$$

where $\mathcal{O}$ is any local observable, and $|(\nu)\rangle$ denotes a state that transforms as the irreducible representation (ν) of H . We can construct a projection operator that projects out a given superselection sector of the Hilbert space. It is

$$P^{(\nu)} = \frac{n_{\nu}}{n_H} \sum_{a \in H} \chi^{(\nu)}(a)^* U(a) , \quad (3.6)$$

where $U(a)$ represents the global H transformation $a \in H$ acting on the Hilbert space. This projection operator can be expressed in terms of the operators $F_{[a]}(\Sigma)$, for it

follows from eq. (3.4) that

$$\frac{F_{[a]}(\Sigma)}{\langle F_{[a]}(\Sigma) \rangle_0} \longrightarrow \frac{1}{n_{[a]}} \sum_{a' \in [a]} U(a') , \quad (3.7)$$

as the surface Σ approaches the surface at spatial infinity.

We can also use the operator $F_{[a]}$ to construct an “Aharonov–Bohm Order Parameter” (ABOP) that probes whether nontrivial superselection sectors actually exist. Let

$$W^{(\nu)}(C) \equiv \chi^{(\nu)} \left[P \exp \left(i \int_C A \cdot dx \right) \right] \quad (3.8)$$

denote the Wilson loop operator in the irreducible representation (ν) . This operator introduces a classical source with charge (ν) propagating on the world line C . The ABOP is defined by

$$A_{[a]}^{(\nu)}(\Sigma, C) \equiv \frac{F_{[a]}(\Sigma) W^{(\nu)}(C)}{\langle F_{[a]}(\Sigma) \rangle_0 \langle W^{(\nu)}(C) \rangle_0} . \quad (3.9)$$

If H quantum numbers can indeed be detected at infinite range, then we expect that

$$\langle A_{[a]}^{(\nu)}(\Sigma, C) \rangle_0 \longrightarrow \frac{1}{n_\nu} \chi^{(\nu)}(a^{k(\Sigma, C)}) , \quad (3.10)$$

in the limit in which Σ and C increase to infinite size, with the closest approach of Σ to C also approaching infinity. Here $k(\Sigma, C)$ is the linking number of the surface Σ and the loop C . (In the abelian case, the ABOP was first described in Ref. 13, and was further elaborated in Ref. 4. The non-abelian generalization was introduced in Ref. 14, and its properties were extensively discussed in Ref. 10.)

The operators $F_{[a]}(\Sigma)$ and $A_{[a]}^{(\nu)}(\Sigma, C)$ can also be constructed in two spatial dimensions. Then Σ becomes a closed curve that can be interpreted as the world line of a vortex–antivortex pair.

4. Charge Transfer

We will now consider the non-abelian Aharonov–Bohm interactions between string loops and point particles, and demonstrate that exchange of H -charge can occur.

The total H -charge of a composite system consisting of a string loop and a charged particle can be measured by studying the Aharonov–Bohm interaction of the composite with other, much larger, string loops. Obviously, then, the total H -charge of the composite must be conserved; it cannot change when the particle winds through the loop. Charge exchange is an inevitable consequence of charge conservation.

To see this, it is convenient to imagine a composite of a string loop and a particle–antiparticle pair, where, initially, both the loop and the pair have zero charge (transform trivially under H). Suppose that the particle transforms as the irreducible representation $D^{(\nu)}$ of H ; the antiparticle transforms as the conjugate representation. Let $\{e_i^{(\nu)} \mid i = 1, 2 \dots n_\nu\}$ denote an orthonormal basis for the vector space on which $D^{(\nu)}$ acts. Then the initial state of the pair has the group–theoretic structure

$$\left| \psi_{\text{in}}^{(\nu)} \right\rangle = \frac{1}{\sqrt{n_\nu}} \left| e_i^{(\nu)*} \otimes e_i^{(\nu)} \right\rangle \quad (4.1)$$

(summed over i). The initial state of the loop is the state $|0; [a]\rangle$ defined in eq. (2.4).

Suppose that the particle and antiparticle are initially at the point x_0 . Then the particle traverses a path C that winds through the string loop and returns to x_0 . After this traversal, the state of the pair and the state of the loop are correlated. The total charge is still zero, but in general the pair and the loop both have a nontrivial charge. We can infer the final charge on the loop by calculating the final charge carried by the pair. In fact, the final charge of the pair is actually independent of the initial charge of the loop; it depends only on the class $[a]$. Thus, to calculate the final charge of the pair, we may take the state of the loop to be the flux eigenstate $|a\rangle$ (where the flux is defined in terms of the path C as in eq. (2.1)). It does not matter how the class representative a is chosen.

Using eq. (3.1), we find that the state of the pair after the traversal is

$$\left| \psi_{\text{fin}}^{(\nu)}, a \right\rangle = \frac{1}{\sqrt{n_\nu}} \left| e_i^{(\nu)*} \otimes e_j^{(\nu)} \right\rangle D_{ji}^{(\nu)}(a) . \quad (4.2)$$

This state $\left| \psi_{\text{fin}}^{(\nu)}, a \right\rangle$ does not transform as a definite irreducible representation of H , but it can, of course, be decomposed into states of definite H -charge. The probability $p^{(\mu)}$ that the H -charge is (μ) can be extracted by using the projection operator $P^{(\mu)}$ defined by eq. (3.6). We find

$$\begin{aligned} p_{\text{pair}}^{(\mu)}(\nu, [a]) &= \left\langle \psi_{\text{fin}}^{(\nu)}, a \left| P^{(\mu)} \right| \psi_{\text{fin}}^{(\nu)}, a \right\rangle \\ &= \frac{n_\mu}{n_\nu n_H} \sum_{b \in H} \chi^{(\mu)}(b^{-1}) D_{nm}^{(\nu)*}(a) \left\langle e_m^{(\nu)*} \otimes e_n^{(\nu)} \left| e_k^{(\nu)*} \otimes e_l^{(\nu)} \right\rangle D_{ki}^{(\nu)*}(b) D_{lj}^{(\nu)}(b) D_{ji}^{(\nu)}(a) \right. \\ &= \frac{n_\mu}{n_\nu n_H} \sum_{b \in H} \chi^{(\mu)}(b^{-1}) \chi^{(\nu)}(bab^{-1}a^{-1}) . \end{aligned} \quad (4.3)$$

As we anticipated, this result is unchanged if a is replaced by $a' \in [a]$.

If the total H -charge is zero, then the composite of string loop and pair has a wavefunction of the form

$$\left| \psi_{[a]}^{(\nu)} \right\rangle = \sum_{\mu} \sqrt{p_{\text{pair}}^{(\mu)}} \left| \text{loop}, \mu^* \right\rangle \otimes \left| \text{pair}, \mu \right\rangle . \quad (4.4)$$

Thus, the probability that the loop carries charge (μ) is given by

$$p_{\text{loop}}^{(\mu)}(\nu, [a]) = p_{\text{pair}}^{(\mu*)}(\nu, [a]) = p_{\text{loop}}^{(\mu*)}(\nu^*, [a]) = p_{\text{loop}}^{(\mu*)}(\nu, [a^{-1}]) . \quad (4.5)$$

Of course, this probability is nonvanishing only if $D^{(\mu)}$ is contained in $D^{(\nu)*} \otimes D^{(\nu)}$ and represents the center of H trivially.

We can directly verify that detectable Cheshire charge now resides on the string loop by studying an appropriate gauge-invariant correlation function. Consider the process depicted in Fig. 2. This process is shown in 2+1 dimensions for ease of

visualization, but the generalization to 3+1 dimensions is straightforward. At time t_1 , a vortex–antivortex pair is created. The flux of the vortex lies in the class $[a]$, and the (initial) H -charge of the vortex pair is trivial. At time t_2 , a particle–antiparticle pair is created. The particle has H -charge (ν) , and the pair is (initially) uncharged. Then the particle winds counterclockwise around the $[a]$ vortex, transferring charge to the vortex pair. Next, another vortex–antivortex pair, with flux lying in the class $[b]$, winds around the (now charged) $[a]$ vortex pair, acquiring an Aharonov–Bohm phase that is sensitive to the charge of the $[a]$ pair. Then the charge- (ν) particle winds clockwise around the $[a]$ vortex, discharging the $[a]$ pair. Finally, the particle–antiparticle pair is annihilated at time t_3 , and the $[a]$ vortex–antivortex pair is annihilated at time t_4 .

If the vortices and charged particles are treated as classical sources, this process is described by the correlation function

$$\left\langle F_{[a]}(C_1) F_{[b]}(C_2) W^{(\nu)}(C_3) \right\rangle_0, \quad (4.6)$$

where C_1 is the world line of the $[a]$ vortex, C_2 is the world line of the $[b]$ vortex, and C_3 is the world line of the charged particle. As shown in Fig. 2, the three loops C_1 , C_2 , and C_3 are joined in a topologically nontrivial configuration known as the “Borromean rings;”^[16] no two loops are linked, yet the loops cannot be separated without crossing.

By considering the case where the loops are large and far apart, and comparing with the case where the loops are unjoined, we can isolate the Aharonov–Bohm factor acquired by the $[b]$ vortex pair that winds around the charged $[a]$ vortex pair. The calculation of eq. (4.6), using weak–coupling perturbation theory on the lattice, is described in Ref. 10. We will not repeat the details of the calculation here, but it is easy to explain the main idea. Loosely speaking, inserting a classical vortex with flux a on the closed path C_1 is equivalent to performing a singular gauge transformation on a surface S_1 that is bounded by C_1 . The path has an orientation, which induces an orientation of the surface. The effect of the singular gauge transformation on the

Wilson loop $W^{(\nu)}(C_3)$ is to insert the factor $D^{(\nu)}(a)$ where C_3 crosses S_1 in a positive sense, and to insert the factor $D^{(\nu)}(a^{-1})$ where C_3 crosses S_1 in a negative sense. In Fig. 3, we see that the loop C_3 successively crosses S_2 in a negative sense, S_1 in a negative sense, S_2 in a positive sense, and S_1 in a positive sense, before closing. Due to the path ordering of the Wilson loop, the factor due to a later crossing appears to the left of the factor due to an earlier crossing. These crossings therefore modify $\langle W^{(\nu)}(C_3) \rangle_0$ by the factor $(1/n_\nu)\chi^{(\nu)}(aba^{-1}b^{-1})$ compared to the case where C_3 is unjoined with C_1 and C_2 . Recalling that a and b are averaged over a class when $F_{[a]}$ and $F_{[b]}$ are inserted, we find that

$$\frac{\langle F_{[a]}(C_1) F_{[b]}(C_2) W^{(\nu)}(C_3) \rangle_0}{\langle F_{[a]}(C_1) \rangle_0 \langle F_{[b]}(C_2) \rangle_0 \langle W^{(\nu)}(C_3) \rangle_0} \longrightarrow \frac{1}{n_H} \sum_{h \in H} \frac{1}{n_\nu} \chi^{(\nu)}(hah^{-1}bha^{-1}h^{-1}b^{-1}) \quad (4.7)$$

when the loops are large, far apart, and joined.

In 3+1 dimensions, there is an analog of the Borromean ring configuration, in which two disjoint closed surfaces Σ_1 and Σ_2 are joined by a closed loop C_3 that does not link with either surface. For this configuration, eq. (4.7) still applies, with C_1 and C_2 replaced by Σ_1 and Σ_2 . We can decompose the right-hand-side of eq. (4.7) into characters as

$$\sum_{\mu} p_{\text{loop}}^{(\mu)}(\nu, [a]) \frac{1}{n_\mu} \chi^{(\mu)}(b) , \quad (4.8)$$

where $p_{\text{loop}}^{(\mu)}(\nu, [a])$ is the probability that the charge carried by the $[a]$ string loop, and detected by the $[b]$ string loop, is (μ) . (Compare eq. (3.4).) Using the orthogonality of the characters, we find from eq. (4.7) and (4.8) that

$$p_{\text{loop}}^{(\mu)}(\nu, [a]) = \frac{n_\mu}{n_\nu n_H} \sum_{b \in H} \chi^{(\mu)}(b^{-1}) \chi^{(\nu)}(aba^{-1}b^{-1}) , \quad (4.9)$$

in agreement with eq. (4.5) and (4.3). Thus, the charge lost by the particle pair has indeed been transferred to the $[a]$ string loop. (Note that, in order to get the right

answer, it is important to choose consistent orientations for the world sheets Σ_1 and Σ_2 —the $[a]$ string must pass through the $[b]$ string in the same sense that the Wilson loop passes through the $[a]$ string. Otherwise, we would in effect be measuring the charge of the $[a]$ string with a $[b^{-1}]$ string, rather than a $[b]$ string.)

We will now derive eq. (4.7) by a different method that invokes the “holonomy interaction” between string loops. Consider two flux-eigenstate string loops that initially carry flux a and b . Now suppose that the b loop sweeps around the a loop and returns to its original position. After this process, the flux of the b loop is unchanged, but the flux of the a loop has been altered; it has become a bab^{-1} loop.^[17,8,6] (Here again, we must be careful about the orientations of the string loops. The a loop becomes a bab^{-1} loop if it passes through the b loop in the same sense as the path C that is used to define the flux of the b loop. If it passes through the b loop in the opposite sense, it becomes a $b^{-1}ab$ loop.)

Return now to the Borromean ring process. Suppose that two string loops are initially in the flux eigenstate $|a, b\rangle$. Then a particle–antiparticle pair is created, and the particle winds through the a loop; the new state of the string loops and the particle–antiparticle pair can be expressed as

$$\frac{1}{\sqrt{n_\nu}} \left| a, b, e_i^{(\nu)*} \otimes e_j^{(\nu)} \right\rangle D_{ji}^{(\nu)}(a) . \quad (4.10)$$

(Compare eq. (4.2).) When the b loop sweeps around the a loop, the state becomes

$$\frac{1}{\sqrt{n_\nu}} \left| bab^{-1}, b, e_i^{(\nu)*} \otimes e_j^{(\nu)} \right\rangle D_{ji}^{(\nu)}(a) , \quad (4.11)$$

due to the holonomy interaction. Now the particle winds back through the bab^{-1} loop (in the opposite sense), and the state becomes

$$\frac{1}{\sqrt{n_\nu}} \left| bab^{-1}, b, e_i^{(\nu)*} \otimes e_k^{(\nu)} \right\rangle D_{kj}^{(\nu)}(ba^{-1}b^{-1}) D_{ji}^{(\nu)}(a) . \quad (4.12)$$

Finally, the particle–antiparticle pair annihilates, and we have

$$\frac{1}{n_\nu} \chi^{(\nu)}(aba^{-1}b^{-1}) |b^{-1}ab, b\rangle . \quad (4.13)$$

To reproduce eq. (4.7), we must take the initial string state to be $|0; [a], 0; [b]\rangle$, in which the $[a]$ and $[b]$ loops are both uncharged. Thus, we average both a and b over a class. We find that the effect of the particle–antiparticle pair on the string state is

$$|0; [a], 0; [b]\rangle \longrightarrow \left(\frac{1}{n_H} \sum_{h \in H} \frac{1}{n_\nu} \chi^{(\nu)}(hah^{-1}bha^{-1}h^{-1}b^{-1}) \right) |0; [a], 0; [b]\rangle . \quad (4.14)$$

By creating the initial string state and annihilating the final string state, we obtain eq. (4.7).

5. A Final Comment

We described in Section 3 how a charge–zero string loop can be used in an Aharonov–Bohm interference experiment to measure the charge of an object. (The corresponding measurement process, using flux eigenstate strings, was described in Ref. 9.) We can imagine doing a double–slit experiment with a beam of particles of unknown charge, where a string loop in the state $|0; [a]\rangle$ surrounds one of the slits. By observing how the shift in the interference pattern depends on the class $[a]$, we can determine the character of the representation according to which the particles in the beam transform, and so infer their charge.

However, the phenomenon of charge transfer raises a puzzle. If a particle passes through the slit that is surrounded by the string, it transfers charge to the string. By measuring the charge on the string loop later, we can find out which slit the particle passed through. Thus, no interference pattern should be seen.

The resolution of this puzzle is that there is a nonvanishing probability, in general, that no charge transfer takes place. This probability is given by eq. (4.9) in the case

where (μ) is the trivial representation (0) ; we then have

$$p_{\text{loop}}^{(0)}(\nu, [a]) = \left| \frac{1}{n_\nu} \chi^{(\nu)}(a) \right|^2. \quad (5.1)$$

Therefore, as long as the character does not vanish, it is possible for the particle to slip through the string loop without being detected, and an interference pattern is observed. From the interference pattern, the phase of the character, as well as its modulus, can be deduced.

We thank Mark Alford, Hoi-Kwong Lo, John March-Russell, and David Wales for useful discussions.

REFERENCES

1. M. Alford, K. Benson, S. Coleman, J. March-Russell, and F. Wilczek, “Interactions and Excitations of Non-Abelian Vortices,” *Phys. Rev. Lett.* **64**, 1632 (1990); **65**, 668 (1990) (E).
2. A.S. Schwarz, “Field Theories With No Local Conservation of Electric Charge,” *Nucl. Phys.* **B208**, 141 (1982); A.S. Schwarz and Y.S. Tyupkin, “Grand Unification and Mirror Particles,” *Nucl. Phys.* **B209**, 427 (1982).
3. P. Ginsparg and S. Coleman, unpublished (1982).
4. J. Preskill and L. Krauss, “Local Discrete Symmetry and Quantum–Mechanical Hair,” *Nucl. Phys.* **B341**, 50 (1990).
5. M. Alford, K. Benson, S. Coleman, J. March-Russell, and F. Wilczek, “Zero Modes of Non-Abelian Vortices,” *Nucl. Phys.* **B349**, 414 (1991).
6. M. Bucher, H.-K. Lo, and J. Preskill, “Topological Approach to Alice Electrodynamics,” Caltech Preprint CALT-68-1752 (1991).
7. M. Alford, J. March-Russell, and F. Wilczek, “Discrete Quantum Hair on Black Holes and the Non-Abelian Aharonov–Bohm Effect,” *Nucl. Phys.* **B337**, 695 (1990).

8. M. Bucher, “The Aharonov–Bohm Effect and Exotic Statistics for Non–Abelian Vortices,” Nucl. Phys. **B350**, 163 (1991).
9. M. Alford, S. Coleman and J. March-Russell, “Disentangling Non–Abelian Discrete Quantum Hair,” Nucl. Phys. **B351**, 735 (1991).
10. M. Alford, K.-M. Lee, J. March-Russell, and J. Preskill, “Quantum Field Theory of Non-Abelian Strings and Vortices,” Caltech preprint CALT-68-1700 (1991).
11. A. Balachandran, F. Lizzi, and V. Rogers, “Topological Symmetry Breakdown in Cholesterics, Nematics, and ^3He ,” Phys. Rev. Lett. **52**, 1818 (1984).
12. L. Krauss and F. Wilczek, “Discrete Gauge Symmetry in Continuum Theories,” Phys. Rev. Lett. **62**, 1221 (1989).
13. K. Fredenhagen, and M. Marcu, “Charged States in Z_2 Gauge Theories,” Comm. Math. Phys. **92**, 81 (1983); K. Fredenhagen, “Particle Structure of Gauge Theories,” in *Fundamental Problems of Gauge Field Theory*, ed. G. Velo and A. S. Wightman (Plenum, New York, 1986).
14. M. Alford and J. March-Russell, “New Order Parameters for Non–Abelian Gauge Theories,” Princeton preprint PUPT-1226 (1990).
15. G. ’t Hooft, “On the Phase Transition Towards Permanent Quark Confinement,” Nucl. Phys. **B138**, 1 (1978); “A Property of Electric and Magnetic Flux in Non-Abelian Gauge Theories,” Nucl. Phys. **B153**, 141 (1979).
16. See, for example, D. Rolfsen, *Knots and Links* (Publish or Perish, Wilmington, 1976).
17. F. Wilczek and Y.-S. Wu, “Space-Time Approach to Holonomy Scattering,” Phys. Rev. Lett. **65**, 13 (1990).

FIGURE CAPTIONS

- 1) The path C , starting and ending at the point x_0 , encircles a loop of string.
- 2) The Borromean rings. C_1 is the world line of an $[a]$ vortex, C_2 is the world line of a $[b]$ vortex, and C_3 is the world line of a charged particle that transforms as the representation (ν) . The charged particle transfers charge to the $[a]$ vortex–antivortex pair, and the charge is subsequently detected via the Aharonov–Bohm interaction of the pair with the $[b]$ vortex.
- 3) A deformation of the rings shown in Fig. 2. The gauge field is singular on the surfaces S_1 and S_2 that are bounded by the loops C_1 and C_2 .